

Determination of weak phases ϕ_2 and ϕ_3 from $B \rightarrow \pi\pi, K\pi$ in the pQCD method

Yong-Yeon Keum

*EKEN Lab. Department of Physics
Nagoya University, Nagoya 464-8602 Japan*

DPNU-02-28

hep-ph/0209002

Presented at CERN CKM Workshop
13–16 February 2002

Abstract

We look at methods to determine the weak phases ϕ_2 and ϕ_3 from $B \rightarrow \pi\pi$ and $K\pi$ decays within the perturbative QCD approach. We obtain quite interesting bounds on ϕ_2 and ϕ_3 from recent experimental measurement in B-factory: $55^\circ \leq \phi_2 \leq 100^\circ$ and $51^\circ \leq \phi_3 \leq 129^\circ$. Specially we predict the possibility of large direct CP violation effect in $B^0 \rightarrow \pi^+\pi^-$ ($23 \pm 7\%$) decay.

1 INTRODUCTION

One of the most exciting aspect of present high energy physics is the exploration of CP violation in B-meson decays, allowing us to overconstrain both sides and three weak phases $\phi_1 (= \beta)$, $\phi_2 (= \alpha)$ and $\phi_3 (= \gamma)$ of the unitarity triangle of the Cabibbo-Kobayashi-Maskawa (CKM) matrix [1] and to check the possibility of New Physics. The “gold-plated” mode $B_d \rightarrow J/\psi K_s$ [2], which allow us to determine ϕ_1 without any hadron uncertainty, was recently measured by BaBar and Belle collaborations [3]: $\phi_1 = (25.5 \pm 4.0)^\circ$. In addition, there are many other interesting channels with which we may achieve this goal by determining ϕ_2 and ϕ_3 [4].

In this letter, we focus on the $B \rightarrow \pi^+\pi^-$ and $K\pi$ processes, providing promising strategies for determining the weak phases of ϕ_2 and ϕ_3 , by using the perturbative QCD method.

The perturbative QCD method (pQCD) has predictive power demonstrated successfully in exclusive 2 body B-meson decays, specially in charmless B-meson decay processes [5]. By introducing parton transverse momenta $k_\perp$, we can generate naturally the Sudakov suppression effect due to resummation of large double logarithms $Exp[-\frac{\alpha_s C_F}{4\pi} \ln^2(\frac{Q^2}{k_\perp^2})]$, which suppress the long-distance contributions in the small $k_\perp$ region and give a sizable average $\langle k_\perp^2 \rangle \sim \bar{\Lambda} M_B$. This can resolve the end point singularity problem and allow the applicability of pQCD to exclusive decays. We found that almost all of the contribution to the exclusive matrix elements come from the integration region where $\alpha_s/\pi < 0.3$ and the perturbative treatment can be justified.

In the pQCD approach, we can predict the contribution of non-factorizable term and annihilation diagram on the same basis as the factorizable one. A folklore for annihilation contributions is that they are negligible compared to W-emission diagrams due to helicity suppression. However the operators $O_{5,6}$ with helicity structure $(S - P)(S + P)$ are not suppressed and give dominant imaginary values, which is the main source of strong phase in the pQCD approach. So we have a large direct CP violation in $B \rightarrow \pi^\pm \pi^\mp, K^\pm \pi^\mp$, since large strong phase comes from the factorized annihilation diagram, which can distinguish pQCD from other models [6, 7].




Fig. 1: Plot of $C_{\pi\pi}$ versus $S_{\pi\pi}$ for various values of ϕ_2 with $\phi_1 = 25.5^\circ$, $0.18 < R_c < 0.30$ and $-41^\circ < \delta < -32^\circ$ in the pQCD method. Here we consider the allowed experimental ranges of BaBar measurement within 90% C.L. Dark areas are allowed regions in the pQCD method for different ϕ_2 values.

2 Extraction of $\phi_2 (= \alpha)$ from $B \rightarrow \pi^+\pi^-$

Even though isospin analysis of $B \rightarrow \pi\pi$ can provide a clean way to determine ϕ_2 , it might be difficult in practice because of the small branching ratio of $B^0 \rightarrow \pi^0\pi^0$. In reality to determine ϕ_2 , we can use the time-dependent rate of $B^0(t) \rightarrow \pi^+\pi^-$ including sizable penguin contributions. In our analysis we use the c-convention. The amplitude can be written as:

$$\begin{aligned} A(B^0 \rightarrow \pi^+\pi^-) &= V_{ub}^* V_{ud} A_u + V_{cb}^* V_{cd} A_c + V_{tb}^* V_{td} A_t, \\ &= V_{ub}^* V_{ud} (A_u - A_t) + V_{cb}^* V_{cd} (A_c - A_t), \\ &= -(|T_c| e^{i\delta_T} e^{i\phi_3} + |P_c| e^{i\delta_P}) \end{aligned} \quad (1)$$

Penguin term carries a different weak phase than the dominant tree amplitude, which leads to generalized form of the time-dependent asymmetry:

$$A(t) \equiv \frac{\Gamma(\bar{B}^0(t) \rightarrow \pi^+\pi^-) - \Gamma(B^0(t) \rightarrow \pi^+\pi^-)}{\Gamma(\bar{B}^0(t) \rightarrow \pi^+\pi^-) + \Gamma(B^0(t) \rightarrow \pi^+\pi^-)} = S_{\pi\pi} \sin(\Delta mt) - C_{\pi\pi} \cos(\Delta mt) \quad (2)$$

where

$$C_{\pi\pi} = \frac{1 - |\lambda_{\pi\pi}|^2}{1 + |\lambda_{\pi\pi}|^2}, \quad S_{\pi\pi} = \frac{2 \operatorname{Im}(\lambda_{\pi\pi})}{1 + |\lambda_{\pi\pi}|^2} \quad (3)$$

satisfies the relation of $C_{\pi\pi}^2 + S_{\pi\pi}^2 \leq 1$. Here

$$\lambda_{\pi\pi} = |\lambda_{\pi\pi}| e^{2i(\phi_2 + \Delta\phi_2)} = e^{2i\phi_2} \left[\frac{1 + R_c e^{i\delta} e^{i\phi_3}}{1 + R_c e^{i\delta} e^{-i\phi_3}} \right] \quad (4)$$

with $R_c = |P_c/T_c|$ and the strong phase difference between penguin and tree amplitudes $\delta = \delta_P - \delta_T$. The time-dependent asymmetry measurement provides two equations for $C_{\pi\pi}$ and $S_{\pi\pi}$ for three (unknown) variables R_c, δ and ϕ_2 .

When we define $R_{\pi\pi} = \overline{Br}(B^0 \rightarrow \pi^+\pi^-)/\overline{Br}(B^0 \rightarrow \pi^+\pi^-)|_{tree}$, where $\overline{Br}$ stands for a branching ratio averaged over B^0 and $\bar{B}^0$, the explicit expression for $S_{\pi\pi}$ and $C_{\pi\pi}$ are given by:

$$R_{\pi\pi} = 1 - 2 R_c \cos\delta \cos(\phi_1 + \phi_2) + R_c^2, \quad (5)$$

$$R_{\pi\pi} S_{\pi\pi} = \sin 2\phi_2 + 2 R_c \sin(\phi_1 - \phi_2) \cos\delta - R_c^2 \sin 2\phi_1, \quad (6)$$

$$R_{\pi\pi} C_{\pi\pi} = 2 R_c \sin(\phi_1 + \phi_2) \sin\delta. \quad (7)$$

If we know R_c and δ , ϕ_2 can be determined from the experimental data on $C_{\pi\pi}$ versus $S_{\pi\pi}$.

Since the pQCD method provides $R_c = 0.23^{+0.07}_{-0.05}$ and $-41^\circ < \delta < -32^\circ$, the allowed range of ϕ_2 at present stage is determined as $55^\circ < \phi_2 < 100^\circ$ as shown in Fig. 1. Since we have a relatively large strong phase in pQCD, in contrast to the QCD-factorization ($\delta \sim 0^\circ$), we predict large direct CP violation effect of $A_{cp}(B^0 \rightarrow \pi^+\pi^-) = (23 \pm 7)\%$ which will be tested by more precise experimental measurement within 3 years. In numerical analysis, since the data by Belle collaboration[8] is located outside allowed physical regions, we only considered the recent BaBar measurement[9] with 90% C.L. interval taking into account the systematic errors:

- $S_{\pi\pi} = 0.02 \pm 0.34 \pm 0.05$ [-0.54, +0.58]
- $C_{\pi\pi} = -0.30 \pm 0.25 \pm 0.04$ [-0.72, +0.12].

The central point of BaBar data corresponds to $\phi_2 = 78^\circ$ in the pQCD method.




Fig. 2: Plot of $\Delta\phi_2$ versus ϕ_2 with $\phi_1 = 25.5^\circ$, $0.18 < R_c < 0.30$ and $-41^\circ < \delta < -32^\circ$ in the pQCD method.

Denoting by $\Delta\phi_2$ the deviation of ϕ_2 due to the penguin contribution, derived from Eq.(4), it can be determined for known values of R_c and δ by using the relation $\phi_3 = 180 - \phi_1 - \phi_2$. In Fig. 2 we show that pQCD prediction on the relation $\Delta\phi_2$ versus ϕ_2 . For allowed regions of $\phi_2 = (55 \sim 100)^\circ$, we have $\Delta\phi_2 = (8 \sim 16)^\circ$. The main uncertainties come from the uncertainty of $|V_{ub}|$. The non-zero value of $\Delta\phi_2$ for $55^\circ < \phi_2 < 100^\circ$ demonstrates sizable penguin contributions in $B^0 \rightarrow \pi^+\pi^-$ decay.

3 Extraction of $\phi_3(=\gamma)$ from $B^0 \rightarrow K^+\pi^-$ and $B^+ \rightarrow K^0\pi^+$ Processes

By using tree-penguin interference in $B^0 \rightarrow K^+\pi^- (\sim T' + P')$ versus $B^+ \rightarrow K^0\pi^+ (\sim P')$, CP-averaged $B \rightarrow K\pi$ branching fraction may lead to non-trivial constraints on the ϕ_3 angle[10]. In order to determine ϕ_3 , we need one more useful information on CP-violating rate differences[11]. Let's introduce the following observables :

$$R_K = \frac{\overline{Br}(B^0 \rightarrow K^+\pi^-) \tau_+}{\overline{Br}(B^+ \rightarrow K^0\pi^+) \tau_0} = 1 - 2 r_K \cos\delta \cos\phi_3 + r_K^2 \geq \sin^2\phi_3 \quad (8)$$

$$\begin{aligned} A_0 &= \frac{\Gamma(\bar{B}^0 \rightarrow K^-\pi^+) - \Gamma(B^0 \rightarrow K^+\pi^-)}{\Gamma(B^- \rightarrow \bar{K}^0\pi^-) + \Gamma(B^+ \rightarrow \bar{K}^0\pi^+)} \\ &= A_{cp}(B^0 \rightarrow K^+\pi^-) R_K = -2r_K \sin\phi_3 \sin\delta. \end{aligned} \quad (9)$$

where $r_K = |T'/P'|$ is the ratio of tree to penguin amplitudes and $\delta = \delta_{T'} - \delta_{P'}$ is the strong phase difference between tree and penguin amplitudes. After eliminate $\sin\delta$ in Eq.(8)-(9), we have

$$R_K = 1 + r_K^2 \pm \sqrt{(4r_K^2 \cos^2\phi_3 - A_0^2 \cot^2\phi_3)}. \quad (10)$$

Here we obtain $r_K = 0.201 \pm 0.037$ from the pQCD analysis[5] and $A_0 = -0.110 \pm 0.065$ by combining recent BaBar measurement on CP asymmetry of $B^0 \rightarrow K^+\pi^-$: $A_{cp}(B^0 \rightarrow K^+\pi^-) = -10.2 \pm 5.0 \pm 1.6\%$ [9] with present world averaged value of $R_K = 1.10 \pm 0.15$ [12].

As shown in Fig.3, we can constrain the allowed ϕ_3 with 1σ range of World Averaged R_K as follows:

- For $\cos\delta > 0$, $r_K = 0.164$: we can exclude $0^\circ \leq \phi_3 \leq 6^\circ$ and $24^\circ \leq \phi_3 \leq 75^\circ$.
- For $\cos\delta > 0$, $r_K = 0.201$: we can exclude $0^\circ \leq \phi_3 \leq 6^\circ$ and $27^\circ \leq \phi_3 \leq 75^\circ$.
- For $\cos\delta > 0$, $r_K = 0.238$: we can exclude $0^\circ \leq \phi_3 \leq 6^\circ$ and $34^\circ \leq \phi_3 \leq 75^\circ$.
- For $\cos\delta < 0$, $r_K = 0.164$: we can exclude $0^\circ \leq \phi_3 \leq 6^\circ$.
- For $\cos\delta < 0$, $r_K = 0.201$: we can exclude $0^\circ \leq \phi_3 \leq 6^\circ$ and $35^\circ \leq \phi_3 \leq 51^\circ$.
- For $\cos\delta < 0$, $r_K = 0.238$: we can exclude $0^\circ \leq \phi_3 \leq 6^\circ$ and $24^\circ \leq \phi_3 \leq 62^\circ$.

From the table 2 of ref.[13], we obtain $\delta_{P'} = 157^\circ$, $\delta_{T'} = 1.4^\circ$, and the negative value of $\cos\delta$: $\cos\delta = -0.91$. The maximum value of the constraint bound for the ϕ_3 strongly depends on the value of $|V_{ub}|$. When we take the central value of $r_K = 0.201$, ϕ_3 is allowed within the ranges of $51^\circ \leq \phi_3 \leq 129^\circ$, which is consistent with the results by the model-independent CKM-fit in the (ρ, η) plane.

4 CONCLUSION

We discussed two methods to determine the weak phases ϕ_2 and ϕ_3 within the pQCD approach through 1) Time-dependent asymmetries in $B^0 \rightarrow \pi^+\pi^-$, 2) $B \rightarrow K\pi$ processes via penguin-




Fig. 3: Plot of R_K versus ϕ_3 with $r_K = 0.164, 0.201$ and 0.238 .

tree interference. We can already obtain interesting bounds on ϕ_2 and ϕ_3 from present experimental measurements. Our predictions within pQCD method is in good agreement with present experimental measurements in charmless B-decays. Specially our pQCD method predicted a large direct CP asymmetry in $B^0 \rightarrow \pi^+ \pi^-$ decay, which will be a crucial touch stone in order to distinguish our approach from others in future precise measurements. More detail works on other methods in $B \rightarrow K\pi$ and $D^{(*)}\pi$ processes will appear elsewhere. [14].

ACKNOWLEDGEMENTS

It is a great pleasure to thank E. Paschos, A.I. Sanda, H.-n. Li, and other members of PQCD working group for fruitful collaborations and joyful discussions. We would like to thank S.J. Brodsky, H.Y. Cheng and M. Kobayashi for their hospitality and encouragement. This work was supported in part by Science Council of R.O.C. under Grant No. NSC-90-2811-M-002 and in part by Grant-in Aid for Scientific Reserach from Ministry of Education, Science and Culture of Japan.

References

- [1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [2] A.B. Carter and A.I. Sanda, Phys. Rev. Lett. **45** (1980) 952; Phys. Rev. **D23**, 1567 (1981); I.I. Bigi and A.I. Sanda, Nucl. Phys. **B193**, 85 (1981); I.I. Bigi and A.I. Sanda, CP Violation, Cambridge University Press, Cambridge, 2000.

- [3] BaBar Collaboration (B. Aubert et al.), Phys. Rev. Lett. **87** (2001) 091801. Belle Collaboration (K. Abe et al.), Phys. Rev. Lett. **87** (2001) 091802.
- [4] Recent works: R. Fleischer and J. Matias, hep-ph/0204101; M. Gronau and J.L. Rosner, Phys.Rev.**D65** (2002) 013004, Erratum-ibid.**D65** (2002) 079901; Phys. Rev. **D65** (2002) 093012; hep-ph/0205323; C.D. Lu and Z. Xiao, hep-ph/0205134.
- [5] Y.Y. Keum, H.n. Li and A.I. Sanda, Phys. Lett. **B504** (2001) 6; Phys. Rev. **D63** (2001) 054008; hep-ph/0201103; Y.Y. Keum and H.n. Li, Phys. Rev. **D63** (2001) 074006; C.D. Lu, K. Ukai and M.Z. Yang, Phys. Rev. **D63** (2001) 074009; C.-H. Chen, Y.Y. Keum and H.-n. Li, Phys.Rev.**D64** (2001) 112002; hep-ph/0204166; S. Mishima, Phys.Lett. **B521** (2001) 252.
- [6] M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, Phys. Rev. Lett. **83** (1999) 1914; Nucl. Phys. **B591**, (2000) 313.
- [7] M. Ciuchini, contributed talk at this workshop.
- [8] Belle Collaboration (K. Abe et al.), Belle-preprint 2002-8 [hep-ex/0204002].
- [9] BaBar Collaboration (B. Aubert et al.), BaBar-Pub-02/09 [hep-ex/0207055].
- [10] R. Fleischer and T. Mannel, Phys. Rev. **D57**, (1998) 2752; M. Neubert and J.L. Rosner, Phys. Lett. **B441** (1998) 403; Phys. Rev. Lett. **81**, (1998) 5076.
- [11] M. Gronau and J.L. Rosner, Phys.Rev.**D65** (2002) 013004, Erratum-ibid.**D65** (2002) 079901.
- [12] R. Bartoldus, talk on Review of rare two-body B decays at FPCP workshop, May 17, 2002.
- [13] Y.Y. Keum, Proceeding at the 3rd workshop on Higher Luminosity B Factory, Aug. 6-7, 2002; hep-ph/0209014; hep-ph/0209208.
- [14] Y.Y. Keum et al., in preparation; Y.Y. Keum, in preparation.